



MMLV Reverse Transcriptase

Applications:

- RT PCR
- Synthesis of cDNA
- mRNA 5'-end Mapping by Primer Extension Analysis
- End-labeling of DNA
- Dideoxynucleotide Sequencing

Description:

MMLV Reverse Transcriptase, encoded by Moloney Murine Leukemia Virus (MMLV RT) is an RNA-dependent DNA polymerase that synthesizes the cDNA first strand from a single-stranded RNA template to which a primer has been hybridized. MMLV RT will also extend primers hybridized to single-stranded DNA. Second strand cDNA synthesis can be achieved from some mRNA templates without an additional DNA polymerase.

Concentration: 200 u/μl

Storage Buffer:

200 mM potassium phosphate (pH 7.2), 0.2% Triton X-100, 2 mM DTT and 50% glycerol

Reaction Buffer 5X:

250 mM Tris-HCl (pH 8.3), 375 mM KCl, 15 mM MgCl₂ and 50 mM DTT

Unit definition:

One unit of the enzyme incorporates 1 nmol dTTP into acid-precipitable material in 10 minutes at 37°C, using poly(A) oligo dT as a template primer.

Quality control:

Endonuclease Activity: 1 μg of Type 1 supercoiled plasmid DNA is incubated with 500 units of enzyme in 1X reaction buffer for one hour at 37°C. The supercoiled DNA is visualized on an ethidium bromide-stained agarose gel to verify absence of nicking or cutting.

Nuclease Activity: 50 ng of radio labelled DNA or RNA is incubated with 200 units of enzyme in 1X reaction buffer for one hour at 37°C, resulting in <1% release for both DNase and RNase.

Purity: >90% as judged by SDS-polyacrylamide gels with blue staining. MMLV RT is free of detectable RNase, and DNase (exo- and endonuclease) activities.

Usage:

Standard Protocol:

We recommend to prepare 2 Mixes

Mix I

Component	Amount/conc.
a. Total RNA or b. PolyA RNA	1-5 μg 50-500 ng
c. Strand-specific primer or d. oligo dT / random primer for each μg of RNA	10 pM 250-500 ng
sterile Water	up to 8 μl
Incubation	Temperature

.. a good decision ..

GeneON .. a good decision ..

Contact Phone +49-(0)-621- 5720 864 Fax: +49-(0)-621-5724 462

E-Mail: <mailto:info@geneon.net> WEB: <http://www.taq-dna.com/> Version: 25.10.2009 AS



10 min	70 °C
10 - 15 min (for c. specific primers) or 5 min (for d. oligo dT / random primer)	room temperature place on ice

Mix II

Component	Amount/conc.
5X reaction buffer	4 µl
dNTP mix (10 mM of each = 40 mM)	1 µl
optional: RNAsin	20-40 units
MMLV Reverse (200 u/µl)	200 units
sterile water	up to 20 µl
combine Mix I and Mix II and gently vortex	

Step	Temperature
30 - 115 min ^{1.)}	37 - 55°C ^{2.)}
10 min (Inactivation of enzyme)	65-70°C

^{1.)} 30 min for cDNA with 500 bp; 115 min for 1,5 kb

^{2.)} depends on the RNA: Higher temperatures (up to 55 °C) for higher structured RNA; Try to adjust the pH to 8.8

Transportation: on blue ice

Storage: at -20°C for 24 months

Ordering information:

Cat.-no	Description	Amount
105-100	MMLV Reverse Transcriptase	10.000 units
105-250	MMLV Reverse Transcriptase	50.000 units

MMLV Reverse Transcriptase

GeneON .. a good decision ..

Contact Phone +49-(0)-621- 5720 864 Fax: +49-(0)-621-5724 462

E-Mail: <mailto:info@geneon.net> WEB: <http://www.taq-dna.com/> Version: 25.10.2009 AS

.. a good decision ..



Cat.-No: 105-100, 10.000 units

Applications:

- RT PCR
- Synthesis of cDNA
- mRNA 5'-end Mapping by Primer Extension Analysis
- End-labeling of DNA
- Dideoxynucleotide Sequencing

Description:

MMLV Reverse Transcriptase, encoded by Moloney Murine Leukemia Virus (MMLV RT) is an RNA-dependent DNA polymerase that synthesizes the cDNA first strand from a single-stranded RNA template to which a primer has been hybridized. MMLV RT will also extend primers hybridized to single-stranded DNA. Second strand cDNA synthesis can be achieved from some mRNA templates without an additional DNA polymerase.

Concentration: 200 u/μl

Storage Buffer:

200 mM potassium phosphate (pH 7.2), 0.2% Triton X-100, 2 mM DTT and 50% glycerol

Reaction Buffer 5X:

250 mM Tris-HCl (pH 8.3), 375 mM KCl, 15 mM MgCl₂ and 50 mM DTT

Unit definition:

One unit of the enzyme incorporates 1 nmol dTTP into acid-precipitable material in 10 minutes at 37°C, using poly(A) oligo dT as a template primer.

Quality control:

Endonuclease Activity: 1 μg of Type 1 supercoiled plasmid DNA is incubated with 500 units of enzyme in 1X reaction buffer for one hour at 37°C. The supercoiled DNA is visualized on an ethidium bromide-stained agarose gel to verify absence of nicking or cutting.

Nuclease Activity: 50 ng of radio labelled DNA or RNA is incubated with 200 units of enzyme in 1X reaction buffer for one hour at 37°C, resulting in <1% release for both DNase and RNase.

Purity: >90% as judged by SDS-polyacrylamide gels with blue staining. MMLV RT is free of detectable RNase, and DNase (exo- and endonuclease) activities.

Usage:

Standard Protocol:

We recommend to prepare 2 Mixes

Mix I

Component	Amount/conc.
a. Total RNA or b. PolyA RNA	1-5 μg 50-500 ng
c. Strand-specific primer or d. oligo dT / random primer for each μg of RNA	10 pM 250-500 ng
sterile Water	up to 8 μl
Incubation	Temperature

.. a good decision ..



10 min	70 °C
10 - 15 min (for c. specific primers) or 5 min (for d. oligo dT / random primer)	room temperature place on ice

Mix II

Component	Amount/conc.
5X reaction buffer	4 µl
dNTP mix (10 mM of each = 40 mM)	1 µl
optional: RNAsin	20-40 units
MMLV Reverse (200 u/µl)	200 units
sterile water	up to 20 µl
combine Mix I and Mix II and gently vortex	

Step	Temperature
30 - 115 min ^{1.)}	37 - 55°C ^{2.)}
10 min (Inactivation of enzyme)	65-70°C

^{1.)} 30 min for cDNA with 500 bp; 115 min for 1,5 kb

^{2.)} depends on the RNA: Higher temperatures (up to 55 °C) for higher structured RNA; Try to adjust the pH to 8.8

Transportation: on blue ice

Storage: at -20°C for 24 months

Ordering information:

Cat.-no	Description	Amount
105-100	MMLV Reverse Transcriptase	10.000 units
105-250	MMLV Reverse Transcriptase	5x10.000 units

.. a good decision ..